



Bayer CropScience LP
 P.O. Box 12014, 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina 27709
 1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

LSP Flowable Fungicide

For The Use On Lentil Seed To Aid In Control Of Seedborne *Ascochyta* Blight.

**SECTION 18 EMERGENCY EXEMPTION
 File Symbol: 06-ND-03**

**FOR DISTRIBUTION AND USE ONLY IN THE STATE OF NORTH DAKOTA
 UNDER SECTION 18 EMERGENCY EXEMPTION**

This labeling must be in possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire Directions for Use and Disclaimer of Warranties before using this product. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product label for LSP Flowable Fungicide (EPA Registration Number 264-948).

Any adverse effects resulting from the use of LSP Flowable Fungicide under this emergency exemption must be immediately reported to the State of North Dakota Department of Agriculture.

EFFECTIVE PERIOD:

This labeling becomes effective March 1, 2006 and expires on June 1, 2006.

DIRECTIONS FOR USE

RECOMMENDED APPLICATIONS		
CROP	DISEASES	RATE OF LSP Flowable
Lentil Seed	<i>Ascochyta</i> Blight	1.7 – 3.0 fl.oz.

Use Directions: To aid in control of seedborne *Ascochyta* Blight: For ready-mix or slurry treaters, mix 1.7 – 3.0 fl.oz. of LSP Flowable Fungicide in an equal amount of water (1:1) to provide adequate coverage and apply to 100 pounds of seed (375 – 750 ppm). Keep dispersion under constant agitation.

A maximum of one application may be made.

Restrictions: When handling thiabendazole, or thiabendazole-treated seed, work in a well-ventilated area and wear chemical resistant gloves; Protective eyewear must be worn when handling thiabendazole; Do not mix with bare hands; Keep out of reach of children; Use good personal hygiene, washing hands and exposed skin before eating, drinking, chewing gum, using tobacco, or using the toilet; no food, drink or tobacco may be allowed in areas of chemical storage or use; May be harmful if swallowed or absorbed through skin; wash exposed skin thoroughly with soap and water after use; Remove clothing immediately if pesticide gets inside; then wash thoroughly and put on clean clothing.

Seed Labeling Requirements

Manual seed treatment is prohibited under this exemption. This product is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Treated seed may be toxic to birds and other wildlife. Clean up any spilled seed immediately, and ensure that treated seed is properly incorporated at planting. Treated seed must not be used for food, feed, or oil processing (left-over treated seed should not be stored but should be double-sown around the headland).

Do not graze or feed livestock on treated areas for four weeks after planting.

Seed Labeling: Treated seed must be labeled in accordance with the U.S. Seed Act as follows:

This seed treated with Thiabendazole at the manufacturer's recommended rate. Do not use for food, feed or oil purposes.

Use with an EPA-approved dye which imparts an un-natural color to the seed.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at www.bayercropscienceus.com

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).