



## TebuStar 3.6L

### For Use on Wheat to suppress *Fusarium* head blight (scab)

SECTION 18 SPECIFIC EXEMPTION

File Symbol: 06-ND-04

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF NORTH DAKOTA  
UNDER SECTION 18 SPECIFIC EXEMPTION

This labeling must be in possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire Directions for Use and Disclaimer of Warranties before using this product. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product label for TebuStar 3.6L (EPA Reg. No. 42750-99).

Any adverse effects resulting from the use of TebuStar 3.6L under this quarantine exemption must be immediately reported to the State of North Dakota Department of Agriculture.

#### EFFECTIVE PERIOD:

This labeling becomes effective 4/14/2006 and expires on 9/01/2006.

#### DIRECTIONS FOR USE

A maximum of 2,000,000 acres of wheat may be treated under this exemption.

#### RECOMMENDED APPLICATIONS

CROP	DISEASES	RATE OF TebuStar 3.6L
Wheat (including durum)	<i>Fusarium</i> head blight (scab)	4 fl oz per acre

#### USE DIRECTIONS

Apply TebuStar 3.6L as a preventive foliar spray when the earliest disease symptoms appear on the leaves or stems. For optimum disease control, the lowest rate of a spray surfactant should be tank-mixed with TebuStar 3.6L. TebuStar 3.6L may be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 3 gallons of spray solution per acre by aircraft spray equipment.

Application through any type of irrigation equipment is prohibited.

A maximum of 1 application per season may be made. A maximum of 4 fl oz of TebuStar 3.6L may be applied per acre per crop season. TebuStar 3.6L may be applied up through the beginning of

flowering (Feekes 10.51). Application may not be made within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. Do not enter treated field for 12 hours after application.

### **ENVIRONMENTAL HAZARDS**

This pesticide is toxic to freshwater, estuarine and marine fish and invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

Do not apply by ground or air within 100 feet of aquatic areas such as lakes, reservoirs, rivers, permanent streams, marshes or natural ponds and estuaries.

For emergencies call CHEMTREC toll free 24 Hours A Day 1-800-424-9300

For PRODUCT USE Information Call 1-800-247-8013 or visit at [www.albaughinc.com](http://www.albaughinc.com)

As with any crop protection product, always read and follow label directions.

Manufactured for:  
Albaugh, Inc.  
1525 NE 36<sup>th</sup> Street  
Ankeny, IA 50021