

2007 Winter Wheat by Fungicide Trial, Prosper, ND

North Dakota State University Experiment Station

Joel Ransom and Scott Meyer

Variety	Yield (bu/A)		Test Weight (lb/bu)	
	No Fungicide†	Fungicide	No Fungicide	Fungicide
Alice	45.7	65.0	56.3	58.3
CDC Buteo	55.0	69.2	58.6	59.8
CDC Falcon	53.9	74.2	55.2	57.8
Darrel	49.4	67.7	55.5	57.7
Expedition	46.8	69.9	56.9	59.2
Goodstreak	57.2	76.7	52.3	55.6
Harding	72.1	84.3	58.4	59.3
Jagalene	31.3	48.3	53.0	57.1
Jerry	67.7	79.7	57.0	58.2
McClintock	48.2	58.8	57.6	58.7
Millennium	76.3	81.9	58.6	59.1
NuDakota	54.1	68.8	53.0	54.9
Paul	31.0	58.3	50.8	55.4
Radiant	38.8	67.9	52.1	57.7
Ransom	65.0	68.5	56.7	57.0
Roughrider	47.9	60.7	58.3	59.7
Wendy	41.9	63.9	55.5	59.0
Wesley	45.6	70.4	53.6	57.1
Yellowstone	35.3	67.5	48.6	54.8
Mean	50.7	68.5	55.1	57.7

† Fungicide treated plots received 3 fl oz/A of Proline™ plus 3 fl oz/A Folicur™ when the varieties were in the early flowering stage.