

2000 Crops Day Report

Pea | Fieldpea Fungicide

2000 Evaluation of POST Applied Fungicides on Field Peas Hettinger

Treatment	Product Rate per Acre		Disease*	1000 Seed Weight	Test Weight	Yield
	1 st Application	2 nd Application	23 DAT			
	June 28	July 10	0-9	grams	lbs/bu	lbs/acre
BASF 500	8.2 oz	0	4.2	221	65.0	2193
BASF 500	5.5 oz	5.5 oz	3.5	209	65.0	2041
Bravo Zn	3 pt	3 pt	3.8	226	64.9	2360
Dithane DF	2 lbs	2 lbs	3.0	212	65.6	2053
Flint	2 oz	2 oz	2.8	208	64.6	2220
Quadris	9.6 oz	0	3.8	220	64.5	2273
Quadris	6.8 oz	6.8 oz	4.0	213	65.3	2073
Untreated	0	0	4.8	218	64.8	2247
Trial Mean			3.7	216	65.0	2183
C.V. %			23.9	3.3	0.9	11.8
LSD 5%			NS	5.2	NS	NS

* Disease rating (23 Days After Treatment): 0 = disease free, 9 = dead plants.

Variety = Majoret

Planting Date: April 28, 2000

Harvest Date: August 1, 2000

NS = no statistical differences between treatments.

Fungicide phytotoxicity was not observed.